

# IPED Field Guide

## Pest Evaluation and Detection



United States  
Department of Agriculture  
Forest Service  
Northeastern Area  
State and Private Forestry  
Newtown Square, PA 19073  
NA-TP-02-10  
May 2010  
[www.na.fs.fed.us](http://www.na.fs.fed.us)



# Credits

---

The IPED Guide Development Team includes professionals from several U.S. Department of Agriculture agencies: Animal and Plant Health Inspection Service; Forest Service, Northern Research Station; Forest Service, Northeastern Area State and Private Forestry, Urban and Community Forestry Program (Donna Murphy); and Forest Service, Northeastern Area State and Private Forestry, Forest Health Protection Program (Daniel Twardus and Joseph O'Brien). Other team members represent the Society of Municipal Arborists, The Davey Institute, University of Georgia Bugwood Network (Joseph LaForest), Cornell University, National Plant Diagnostic Network, and Purdue University.

The cover photo in the lower left was taken by David Bloniarz, U.S. Forest Service, Northern Research Station. The other two photos were taken by Donna Murphy, U.S. Forest Service, Northeastern Area State and Private Forestry.

# IPED Field Guide

## Pest Evaluation and Detection





United States  
Department of Agriculture  
Forest Service  
Northeastern Area  
State and Private Forestry  
11 Campus Blvd., Suite 200  
Newtown Square, PA 19073  
May 2010  
[www.na.fs.fed.us](http://www.na.fs.fed.us)



# Scales

---



[www.vendian.org/mncharity/dir3/paper\\_rulers/](http://www.vendian.org/mncharity/dir3/paper_rulers/)

## Table of Contents

Introduction.....	1	Salt/Chemicals .....	16
Background.....	1	<b>Signs and Symptoms of Foliage and</b>	
Rationale .....	2	<b>Twigs .....</b>	<b>18</b>
<b>Tree Stress.....</b>	<b>3</b>	Defoliation.....	18
Dieback.....	3	Defoliation, >10% of foliage, but not	
Twig Dieback in upper/outer crown		pervasive.....	18
(>10%), but not pervasive .....	4	Defoliation, pervasive throughout	
Pervasive Twig Dieback throughout		the crown .....	20
the Crown .....	4	Leaf Mining, >10% of foliage,	
Epicormic Sprouts .....	4	but not pervasive.....	20
Wilted Foliage .....	6	Leaf Mining, pervasive throughout	
Wilt, Whole Crown .....	6	the crown .....	20
Wilt, Partial Crown .....	6	Chewing of the Midrib Only.....	22
Environmental Stress.....	8	Discolored Foliage >10% Threshold.....	22
Frost Cracks .....	8	Mottling, Spots, or Blotches .....	22
Lightning Strike .....	8	Marginal Scorching (browning) of	
Hail Injury.....	8	Leaves.....	22
Sunscald .....	10	Interveinal Scorching (browning)	
Broken Branches .....	10	of Leaves .....	24
Flooding .....	10	White Coating .....	24
Drought/Poor Soil .....	12	Black Coating (often sticky) .....	24
Human-Caused Stress.....	12	Complete Browning/Bronzing of	
Topping/Poor Pruning .....	14	Leaves or Needles.....	26
Poor or Restricted Planting/ Mulching .....	14	Complete Yellowing of Leaves or	
Wounding of Woody Tissues .....	16	Needles .....	26
		Stippling.....	26

Yellow/Orange/White Pustules .....	26	Shot Holes (<2 mm).....	38
Abnormal Foliage.....	28	Other Holes .....	38
Foliage/Twigs Distorted (including		Insect Presence .....	40
galls on foliage).....	28	Caterpillars .....	40
Witches' Brooms Present .....	28	Beetles.....	40
Insect Signs.....	30	Aphids .....	40
Caterpillars/Sawflies feeding		Scale.....	40
throughout the Crown.....	30	Carpenter Ants .....	42
Beetles feeding throughout the Crown	30	Disease Signs.....	42
Aphids/White Cotton pervasive		Decay .....	42
throughout the Crown.....	30	Conks .....	44
Bags pervasive throughout the Crown	32	Fleshy Mushrooms.....	44
Scale pervasive throughout the		Cankers .....	44
Crown .....	32	Bleeding/Slime Flux .....	44
Tents/Webbing on more than One		Resinosis/Gummosis.....	46
Branch .....	32	Woody Galls or Burls.....	46
Percentage Foliage Affected.....	34	Problem Location .....	46
>10% ; < 30% of the Crown Affected	34	Branches.....	46
>30% but not the Whole Crown .....	34	Bole and/or Root Collar .....	46
Whole Crown Affected .....	34	Loose Bark .....	48
<b>Signs and Symptoms of Branches</b>		Loose Bark .....	48
<b>and Bole.....</b>	<b>36</b>	Rhizomorphs Present .....	48
Insect Signs.....	36	Mycelial Fans or Pads Present .....	48
Frass Only .....	36	Insect Boring or Galleries causing	
Sawdust.....	36	Loose Bark .....	48
Pitch/Resin Exudation.....	36	Glossary.....	51
D-Shaped Exit Holes.....	38	Other Resources .....	57
Pencil Round or Oval Exit Holes		Photo Credits .....	58
(>=2 mm).....	38		

# Introduction

---

The Inventory Pest Evaluation and Detection (IPED) protocol provides a portable, accessible, and standardized method of observing a tree for possible insect or disease problems. It is intended to be a standardized protocol for long-term urban pest detection and monitoring throughout the United States.

This field guide will help you identify the signs and symptoms of tree stress, insect pests, and diseases, which will enable you to make informed, systematic decisions when collecting data using the IPED protocol. This guide is not intended to be used for diagnostic work, but rather as a resource guide for field use that corresponds with data entry fields for the IPED desktop and IPED PDA application found within the i-Tree Streets application at: <http://www.itreetools.org/>. An expanded version of this guide can be found online at: <http://wiki.bugwood.org/IPED>.

## Background

---

Urban areas are frequently the first site of introduction for exotic pests, where they remain undetected until populations are well established and have harmful impacts on the health of host trees. Many communities routinely complete and update tree inventories, but often overlook pest invasions because pest detection tools are not part of the inventory process. Prior to the development of IPED, there were no processes in place for aggregating pest inventory data into a standardized form, which would allow communities to analyze pest trends otherwise difficult to detect across geographic or political boundaries.

Urban tree inventories are conducted by communities to better understand the structure, function, and management needs of their urban forests. Detecting tree pest problems during an inventory provides information that can lead to preventative actions that can save trees.

IPED is a tool for integrating field data collection and desktop reporting within the i-Tree suite of tools. i-Tree was created to develop, disseminate, support, and refine urban forest analysis tools and is utilized as the platform for IPED within the i-Tree v3.0 Streets application. For more information, see the field data collection procedures and desktop operation guide within the i-Tree Streets application at [www.i-treetools.org](http://www.i-treetools.org).

## Rationale

---

Since most exotic pests are initially introduced into urban areas (Asian longhorned beetle, emerald ash borer, and gypsy moth, for example), incorporating exotic pest detection into street tree inventories expands detection capabilities and enables communities to become a front line in the detection of exotic pests. IPED provides a portable, standardized method of observing a tree for signs and symptoms of possible insect or disease problems. Awareness of signs and symptoms that indicate potential tree health issues offers early opportunities for preventative actions and improved urban tree management. The rationale behind the field guide builds on the pretense that diagnosis is based largely upon determining signs and symptoms to narrow the range of possible causes of tree health issues such as stress, pests, or disease.

The protocol of using signs and symptoms when observing the foliage, twigs, branches, and boles of trees provides a methodical, systematic approach to pest detection. For the purpose of this field guide, *a symptom is defined as an injury by—or a plant's response to—a pest agent. A sign is defined as the pest organism itself, its skeleton, or a product produced by the pest. A sign is helpful in identifying the cause of a symptom.*

# Tree Stress

---

Tree stress can result from mechanical, physical, or biological factors, or a combination of these. Factors that cause stress can be biotic or abiotic. Stress responses vary by cause and can include slow growth; sparse, undersized, or distorted, often *chlorotic*, nutrient-deficient leaves; browning of leaf margins; premature autumn color; premature leaf drop; and *adventitious sprouts*. Two general sequences of symptoms are recognized. If decline is initiated by an event such as root cutting or severe defoliation, buds and twigs may die as a result of shock. If decline results from chronic stress, foliar symptoms and slow growth are likely to precede dieback.

## Dieback

Crown dieback is the dying back of branches and branch tips generally in the upper and outer portions of the tree crown. Dieback often occurs as a sign of stress but may be associated with a new pest or disease. Crown dieback can be caused by a variety of factors including insects, disease, and environmental conditions. Branches that are defoliated by insects are not considered dieback. When conifers die back due to stress, they tend to lose needles from the inside of the crown out and from the bottom up.

Dieback of twigs or whole branches is often a symptom of either *biotic* or *abiotic stress*, including drought, chemical injury, root injury, insects, or disease.

If you know that the observed dieback is caused by drought or other stresses, record it in the Tree Stress category. Otherwise it should be recorded under foliage symptoms.

# Tree Stress

---

## Dieback

### **Twig Dieback in upper/outer crown (> 10%), but not pervasive**

Dieback of twigs in the upper/outer crown (>10%) can be an early indication of more severe problems such as physical damage to tree roots, root girdling, pollution, drought, soil compaction, or poor drainage. Although 10% dieback of the outer crown may not be detrimental to the health of a tree, it should be considered seriously and warrants a full health assessment. When conifers die back due to stress, they tend to shed needles from the inside of the crown out and from the bottom up.

### **Pervasive Twig Dieback throughout the Crown**

*Pervasive twig dieback* throughout the crown is an indication that the health of the tree is compromised. Pervasive dieback can be a result of long-term drought stress, soil compaction, nutrient deficiencies, poor drainage, girdling roots, insect infestation, or disease.

## Epicormic Sprouts

*Epicormic sprouts*, whether at the base of the tree, along the bole, or on branches, can be caused by topping, wounding, insect infestation, or disease. Epicormic sprouts are weakly attached to the stem. Weak branch attachments will eventually result in branches that are more prone to breakage and decay.



Twig Dieback in upper/outer crown (> 10%), but not pervasive—Maple in urban environment



Pervasive Twig Dieback throughout the crown



Pervasive Twig Dieback throughout the crown—Spruce decline



Epicormic Sprouts—Developing on ash after emerald ash borer attack

# Tree Stress

---

## Wilted Foliage

**If you know that drought or other stress factors are responsible for the wilt observed, record it here. Otherwise it should be recorded under foliage symptoms.**

Wilt is caused by the loss of water pressure in a leaf, causing it to droop, curl, or lose its normal color. Wilt can be caused by insects, disease, or environmental factors such as drought, chemical damage, girdling, root damage, soil compaction, or construction damage. The pest may be located on the branch, twigs, or trunk depending on the time of day the inspection is conducted. Several diseases, such as oak wilt and mimosa wilt, cause foliage wilting. Dutch elm disease causes flagging, which is the wilting and dying of foliage on entire branches.

## Wilt, Whole Crown

Wilting of an entire crown can be caused by abiotic or biotic factors. Abiotic causes are often associated with structural damage to the tree, water deficiency, and/or soil compaction. Biotic factors are often a cause of vascular disease. Vascular wilts are caused by *pathogens* that establish in the xylem and prevent water movement throughout the tree. Vascular wilts are usually fatal, although in some cases symptoms can be limited to branch mortality.

## Wilt, Partial Crown

Partial wilting of the crown can be caused by abiotic or biotic factors. Abiotic causes are often associated with structural damage to the tree, water deficiency, and/or soil compaction. Biotic factors are often associated with pathogens that establish in the xylem. These pathogens prevent nutrients and moisture from moving throughout the tree. Vascular wilts are usually fatal although in some cases symptoms can be limited to branch mortality.




Wilt, Whole Crown—Oak wilt



Wilt, Whole Crown—Heavy wilt symptoms on oak



Wilt, Whole Crown—Pine wilt nematode



Wilt, Partial Crown—American elm



Wilt, Partial Crown—Crown wilt caused by *Verticillium* wilt

# Tree Stress

---

## Environmental Stress

Environmental stress contributes to many problems associated with tree health. Stressful environmental conditions such as extreme temperatures; poor soil; and physical damage to leaves, bark, and roots can predispose a tree to secondary insect and disease attacks.

### Frost Cracks

Frost cracks are caused by differential expansion and contraction of woody stem tissues during extremely cold periods. Trees with thin bark are more prone to frost cracking. These cracks are usually found on the lower bole of the tree and are often associated with other tree defects such as cankers or pruning wounds. Other openings in trees can be caused by decay organisms that cause loss of wood by their actions.

### Lightning Strike

Lightning injury is influenced by a tree's location, moisture content, and the amount of moisture in the soil around the tree. Damage can be correlated to where the tree is struck and the concentration of water in the tree. If the lightning strike is concentrated in the *phloem* between the bark and the wood, then the strike will follow this channel and create an explosive separation of the bark. If there is more moisture in the center of the tree, the explosion from within may blow the tree apart.

### Hail Injury

Young trees exposed to hail typically develop small to large multiple wounds on the exposed side of the tree, damaging the bole and twigs, stripping the bark, and destroying the buds. If the number and size of the wound areas are too large, the tree may die or begin to decline. Older conifers and hardwoods typically have thicker bark, thus they typically suffer less severe damage.



Frost Cracks—Frost and winter injury on the bole



Lightning Strike—Oak



Lightning Strike—Damage on bole of oak



Hail Injury—Hail damage on bark of young peach tree



Hail Injury—Hail damage, cankering on shortleaf pine

# Tree Stress

---

## Environmental Stress

### Sunscald

A variety of cultural practices, such as deep planting and trunk and root injuries, are suspected of being *causal factors* in the development of sunscald. A species characteristic thought to play a role in the development of sunscald is bark thickness. Young, thin-barked species appear to be more susceptible to sunscald but become less susceptible as they mature and develop thicker bark.

### Broken Branches

Trees should have a strong trunk with sturdy, well-spaced branches. The strength of the trunk/branch union depends on the relative sizes of the trunk/branches, the trunk/branch angles, and the spacing of the branches. Broken branches pose serious health risks to trees, contribute to property damage loss, and impact personal safety.

### Flooding

Flooding can severely impact tree health. Injuries related to flooding can contribute to the introduction of diseases in tree roots and at injury sites. It can compromise the stability of a tree's root system and cause stress, resulting in tree decline and/or insect infestations and diseases.



Sunscald—Sunscald damage on one side of young tree



Broken Branches—Broken branch on pine



Flooding—Flooding-related symptoms

# Tree Stress

---

## Environmental Stress

### Drought/Poor Soil

**If you know that drought or poor soil is responsible for the symptoms, record it here. Otherwise it should be recorded under Signs and Symptoms of Foliage and Twigs, Subcategory Discolored Foliage (see page 22).**

Drought contributes to tree stress, making trees more prone to insect and disease attacks. Drought stress can decrease resistance to trunk-invading organisms such as wood borers, fungi, and vascular diseases. Drought also limits nutrient availability by preventing the absorption of essential nutrients. A soil can be adequate for proper plant growth but compromised through compaction or poor drainage. Poor soil contributes to tree stress and manifests itself in many ways, including chlorosis/discoloration of leaves, stunted growth, or gradual dieback of the tree from the top down.

## Human-Caused Stress

Human activities such as poor pruning, improper handling during transplanting, improper planting, construction damage, and the use of *deicing salts* or *herbicides* can damage trees directly or indirectly. Damaged trees can die suddenly or deteriorate over a period of time and are often at increased risk of failure as a result of wounding that does not heal properly. Trees with human-caused stresses are more susceptible to secondary damage-causing agents.



Drought/Poor Soil—Scorching of leaves due to drought damage on sugar maple



Drought/Poor Soil—Drought damage on pine



Drought/Poor Soil—Iron deficiency on oak



Drought/Poor Soil—Magnesium deficiency on leaf

# Tree Stress

---

## Human-Caused Stress

### **Topping/Poor Pruning**

Topping is a pruning practice that harms trees and should not be used. Topping—the pruning of large upright branches between nodes—is sometimes done to reduce the height of a tree. Topping results in epicormic sprouting and contributes to branch death to the next lateral branch below. Poor pruning practices such as tipping, bark cutting, and flush cutting result in bark ripping and the dying back of branch stubs. Poor pruning also provides entry points for canker fungi that kill the cambium, and prevents or delays woundwood formation.

### **Poor or Restricted Planting/Mulching**

Poor or restricted planting space can have detrimental impacts on tree health. Restricted planting space contributes to nutrient and water deficiencies caused by inadequate soil volume. It can also result in stress related to poor drainage; compromised root systems due to the inability of roots to penetrate compacted soil; girdling and encircling roots; and damage to sidewalks, pavement, and utility infrastructure.

Proper mulching of trees can provide organic material, prevent weed establishment, prevent lawn mower damage, and conserve water. Improper mulching techniques such as using noncomposted mulch can rob a soil of its nitrogen or induce nitrate-related burning of feeder roots.

Volcano mulching, the piling of mulch in a high mound at the base of a tree, can result in girdling of the tree due to the decay of bark where it is in contact with the mulch. Poor planting techniques often result in girdling roots if the roots of trees are not released from circular growing patterns at the time of planting, or the soil is too richly amended within the planting hole, thus contributing to circular growing roots that can become girdling roots over time.



Topping/Poor Pruning—Topping-caused epicormic branching



Topping/Poor Pruning—Flush cuts prevent the formation of woundwood



Poor or Restricted Planting—Decline



Poor or Restricted Planting—Decline



Poor Mulching—Improper volcano mulching

# Tree Stress

---

## Human-Caused Stress

### Wounding of Woody Tissues

Trees can easily heal small wounds. In urban areas, tree wounding is often caused by human activity related to the use of machinery, automobiles, lawn mowers, weedwackers, and vandalism that is repeated again and again on the same part of the tree. This constant wounding prevents a tree from fully healing wounds and is an entry point for pests and disease.

### Salt/Chemicals

Salt injury occurs when salt is deposited by spray or *drifts* onto dormant stems and buds of deciduous woody plants and on the stems, buds, and leaves of evergreens. Injury may also occur when excessive amounts of salt accumulate in the root zone of plants. This buildup occurs along city streets, driveways, and sidewalks when salt runoff washes into the soil and when snow laden with salt is plowed and shoveled onto boulevards and lawns.

Chemical damage occurs when drift and misapplication of herbicides damage *nontarget* trees. The total extent of such damage remains unknown, but localized severe damage can occur. The causal agent is often identified through *symptom* expression and by determining the method and rate of nearby herbicide applications. Symptom expression can be variable for a given chemical and is often unreliable when used as the only diagnostic tool.



Wounding of Woody Tissues—Mechanical wound with resin



Wounding of Woody Tissues—Human activity



Salt Damage—Salt injury on aspen



Chemical Damage—Herbicide damage on dogwood

# Signs and Symptoms of Foliage and Twigs

---

For most signs and symptoms observed, a threshold level of 10% should be exceeded prior to noting it in the IPED protocol. For example, a single caterpillar, with no evidence of defoliation or other injury, should not be reported. Symptoms on leaves, twigs, or branches not affecting >10% of the crown should not be reported. Minor mechanical injury on the stem and branches should not be reported.

Observe the foliage and twigs throughout the crown for signs of insect feeding, discoloration of the foliage, or abnormal growth of the foliage or twigs. Observing the exact nature of the feeding is often important in diagnosing the cause of the feeding even if no insect is present. For discolored foliage, the exact nature of the discoloration is also important.

## Defoliation

Foliage eaten wholly or partially is generally referred to as defoliation. The extent of defoliation is often related to the severity of infestation. Foliage feeding can be recognized by the absence of foliage or from uneaten leaf or needle parts. Many insects feed by eating only the softer parts of leaves, resulting in a skeletonizing effect. Leaf miners bore inside the foliage and eat the tissues between the foliage surfaces, resulting in serpentine or blotchy areas. The adult Asian longhorned beetle has the somewhat unusual habit of sporadic feeding of the midrib portions of host leaves. Signs of this feeding may be detected through careful observation of foliage on potential host trees including maples, horse chestnut, birch, elm, poplar, and willow.

### **Defoliation, >10% of foliage, but not pervasive**

Defoliation from any amount of feeding resulting in damage on 10% or more of the leaves/needles within a tree's crown meets the >10% defoliation threshold and should be noted in the IPED protocol.



Defoliation, >10%, but not pervasive—  
European hornbeam



Defoliation, >10%—Defoliation threshold  
must be determined by looking at the  
entire crown



Defoliation, sawfly defoliation >10%  
but not pervasive



Defoliation—If 10% or more of the  
leaves in the tree's crown look like this,  
it meets the >10% defoliation threshold

# Signs and Symptoms of Foliage and Twigs

---

## **Defoliation**

### **Defoliation, pervasive throughout the crown**

Pervasive thinning of leaves in tree crowns is an indication of severe insect infestation and can result in severe stress. Signs of insects causing defoliation may or may not be present, so it is important to be aware of symptoms resulting from the presence of defoliating insects or diseases. It is important to look carefully and determine the feeding pattern of the insect on remaining leaves.

Trees commonly survive the first year of defoliation, although defoliation for one growing season can have an impact on future tree growth. Defoliated trees can grow new leaves later in the growing season, but the leaves are often stunted in appearance. Defoliation that reoccurs can cause tree death. Forest tent caterpillars and Eastern tent caterpillars often defoliate trees entirely, but leave telltale signs such as webbing and tents.

### **Leaf Mining, >10% of foliage, but not pervasive**

Leaf mining is often recognized by the serpentine pattern of feeding beneath the leaf surface as insects live or feed within a leaf. Make sure the threshold of 10% is reached before noting it in the IPED protocol.

### **Leaf Mining, pervasive throughout the crown**

Leaf mining is often recognized by the serpentine pattern of feeding beneath the leaf surface. The locust leafminer and elm leafminer are common pests throughout Eastern North America. Leafminer larvae tunnel through the leaf forming blotches and discoloration. As the miners move to the outer edge of the leaf, the leaf turns brown.



Defoliation, pervasive—Gypsy moth



Leaf Mining, >10% of foliage, but not pervasive—If 10% or more of all needles look like this, it meets the >10% defoliation threshold



Leaf Mining, >10% of foliage, not pervasive—Birch leafminer



Leaf Mining, pervasive throughout the crown—Elm leafminer

# Signs and Symptoms of Foliage and Twigs

---

## Defoliation

### **Chewing of the Midrib Only (any level)**

Feeding along the *midrib* of host foliage can easily be missed unless carefully observed. The Asian longhorned beetle (ALB) will feed on the midrib. Pay particular attention to maples and other ALB hosts for this sign of insect feeding. If midrib feeding is observed, it must be immediately noted in the IPED protocol even if it does not meet the >10% defoliation threshold. It is important to then check for other signs or symptoms of ALB such as frass, exit holes, or dieback.

### **Discolored Foliage (>10% threshold)**

Discolored foliage can be caused by pathogens, insect activity, nutrient deficiency, air pollution, disease, or natural processes such as leaf senescence. Careful observation of the pattern and location of discoloration on the leaves is key to identifying leaf problems. In most cases a 10% defoliation must be met for the IPED protocol.

### **Mottling, Spots, or Blotches**

Mottling is spotting of leaves or needles in an uneven and undefined pattern, usually involving several different shades of color. Leaf spots are more well defined while blotches are usually larger and more irregular in form.

### **Marginal Scorching (browning) of Leaves**

Burn-like damage to leaf margins can be the result of environmental stress or disease. *Marginal scorching* of leaves may be caused by pathogens, drought stress, nutrient deficiency, and some feeding insects. Oak wilt and bacterial leaf scorch cause marginal scorching in leaves.



Defoliation, ALB Chewing of the Midrib Only—Does not need to meet 10% defoliation threshold



Discolored Foliage, Mottling, Spots, or Blotches—Bacterial leaf spot symptoms meet 10% Defoliation threshold



Discolored Foliage, Marginal Scorching—Oak wilt



Discolored Foliage, Marginal Scorching—Dogwood anthracnose

# Signs and Symptoms of Foliage and Twigs

---

## Discolored Foliage

### Interveinal Scorching (browning) of Leaves

*Interveinal scorching* is represented by discoloration in the tissues between the leaf veins. The veins themselves may remain green for some time, and are the last parts of the leaves to discolor. Interveinal chlorosis or scorching is often caused by air pollutants or chemical injury.

### White Coating

White coatings on leaves can arise from the actions of both fungi and insects. Notable among fungi are the powdery mildew pathogens that produce a white material on the surface of leaves. This material on the leaf surface is actually composed of *fungus tissue*, but the fungus also grows into the interior of the leaves, disrupting their ability to photosynthesize. Anthracnose diseases can also sometimes cause a white discoloration on the surface of *necrotic leaf blotches*. Insects such as scales and adelgids often produce a white, cottony protective covering during the stage of their life cycles when they are immobile.

### Black Coating (often sticky)

Sooty molds are caused by several fungi that grow on the surface of leaves, needles, bark, or even wood or concrete structures. The fungi grow on the exudations of honeydew-secreting, sucking insects, such as aphids. Honeydew is a sticky, sweetish substance that is a good substrate for the growth of several different fungi that have very dark mycelia. Although the fungi are not pathogenic to plants, they can cause stress by coating leaves with a heavy black coating and disrupting photosynthesis.



Interveinal Scorching—Ozone injury on maple



Interveinal Scorching—Bacterial leaf scorch



White Coating—Powdery mildew on oak



Black Coating (often sticky)—Sooty mold on rhododendron

# Signs and Symptoms of Foliage and Twigs

---

## Discolored Foliage

### Complete Browning/Bronzing of Leaves or Needles

Browning or bronzing of foliage can be caused by insects, disease, or environmental conditions. An insect or disease relationship may not be obvious in the foliage. Several diseases, such as oak wilt and bronze leaf disease of aspen, cause foliage bronzing. Bronzing can also be caused by some types of mites, ozone damage, and/or pollution.

### Complete Yellowing of Leaves or Needles

Complete yellowing of the leaves is called chlorosis. Chlorosis can be caused by insects, disease, nutrient imbalances, or environmental conditions. An insect or disease relationship may not be obvious in the foliage. If a pest is present, it may be located on the branches, twigs, or trunk. Complete yellowing of all foliage on a tree is often an indication of a problem in the root system. Conifers affected by the root disease pathogen *Armillaria*, or by root collar weevils, often turn completely yellow-green to yellow before they die.

### Stippling

**Stippling** is small, white-spotted, speckled injuries on foliage caused by certain insects, mites, and viruses. In some cases it can be yellow or black, but it is generally white.

### Yellow/Orange/White Pustules

Yellow or orange-yellow, orange, or white spots on foliage are often caused by rust diseases. The pustules can range in color from off-white to orange. Both conifers and hardwoods are susceptible to a variety of rust diseases. Symptoms can occur from spring into fall, depending upon the disease.



Complete Browning/Bronzing of Leaves—Dutch elm disease




Complete Browning/Bronzing of Needles—Hemlock needle blight



Complete Yellowing of Leaves—Iron deficiency on oak



Stippling—Lacebugs



Yellow/Orange Pustules—Eastern gall rust on oak host

# Signs and Symptoms of Foliage and Twigs

---

## **Abnormal Foliage**

There are several foliage symptoms that do not involve discoloration or chewing. These include surface molds, distorted or curled foliage or twigs, and galls.

### **Foliage/Twigs Distorted (including galls on foliage)**

Several insects and diseases can distort foliage and/or twigs. Distortion or stunting can be caused by aphids, shoot moths, cone worms, fungal pathogens, and bacteria. Distorted foliage can also result from herbicide injury. *Leaf rollers* are insects that construct shelters by rolling leaves with silk webbing. These rolled leaves may give the appearance of distorted foliage.

Several insects and diseases can cause shoot tips to curl. Curled tips develop during leaf or needle expansion due to a disruption in the physiology of the tissue. White pine weevil, spruce gall adelgid, and shoot blights caused by fungi commonly cause curling of shoot tips. Fire blight will also cause tip curling. This is referred to as Shepherd's Crooking.

### **Witches' Brooms Present**

*Witches' brooms* are deformities in a tree that change the natural structure of branches or twigs. Witches' brooms often arise as the result of infection by mistletoes, fungi, insects, or disease.



Foliage Distorted—Eastern gall damage



Foliage Distorted—Cherry scallop shell moth



Galls on Foliage—Cooley spruce gall



Witches' Brooms—Hackberry

# Signs and Symptoms of Foliage and Twigs

---

## **Insect Signs**

Observe the foliage and twigs for the presence of insects. It is possible that the signs of insect feeding are present but the insect cannot be found. Some signs of insect presence include cottony masses created by aphids, small bags of foliage created by an insect within which it feeds, or webbing and/or tents.

### **Caterpillars/Sawflies feeding throughout the Crown**

Caterpillars are mostly found on the foliage where they feed. However, occasionally caterpillars can be found within the branches even though they are feeding elsewhere on foliage. When caterpillars are found on twigs, shoots, or branches, look for signs of defoliation. Sawflies belong to the same group of insects as the wasps, ants, and bees. Sawfly larvae look like caterpillars. Some sawfly larvae curl up when touched and some feed in clusters.

### **Beetles feeding throughout the Crown**

Not all beetles feed as adults, but some do, such as the elm leaf beetle. Asian longhorned and emerald ash borer beetles may be observed on the foliage or twigs even though they are not feeding.

### **Aphids/White Cotton pervasive throughout the Crown**

*Aphids* are sap sucking insects. Severe infestations result when aphids can be found throughout the entire crown on twigs and shoots or young branches. Look for aphids on the twigs between the foliage. Some aphids, such as the hemlock woolly adelgid, cover themselves in a white cottony mass for protection. Sometimes aphids are discovered by the appearance of the foliage.



Sawflies—Larch sawfly



Beetles—Asian longhorned beetle adult



Beetles—Emerald ash borer adult



Aphids/White Cotton—Woolly adelgid

# Signs and Symptoms of Foliage and Twigs

---

## **Insect Signs**

### **Bags pervasive throughout the Crown**

Small *spindle-shaped bags* hanging from the foliage or twigs are the protective feeding enclosures for some caterpillars. The bags are constructed from foliage and twigs. Bagworms are common on juniper, arborvitae, spruce, pine, and cedar. Severe infestations result when bagworms can be found throughout the entire tree.

### **Scale pervasive throughout the Crown**

Scale insects often secrete a scale-like wax coating over their backs and can resemble bark. Most scales infest the twigs and smaller branches. The pine needle scale is a serious pest of ornamental pines. These scales can be found on the needles, especially of Scots and mugo pine. Severe infestations result when scales are found throughout the entire tree.

### **Tents/Webbing on more than One Branch**

Silken tents are sometimes created as a protective barrier that an insect will feed in or use as a shelter. The Eastern tent caterpillar is a common example, as are the fall webworm and the uglynest caterpillar.



Bags pervasive throughout the crown—Bagworm



Scales—Hemlock scale



Scale—Soft scale



Tents/Webbing—Satin moth



Tents/Webbing—Fall webworm



Tents/Webbing—Eastern tent caterpillars

# Signs and Symptoms of Foliage and Twigs

---

## **Percentage Foliage Affected**

Considering the whole crown, how much of the foliage is affected by foliage feeding, discoloration, or abnormal growth?

### **>10% ;< 30% of the Crown Affected**

>10-30% of the foliage or crown is affected by feeding, discoloration, or abnormal growth.

### **>30% but not the Whole Crown**

More than 30% of the foliage or crown is affected but not the entire tree.

## **Whole Crown Affected**

Defoliation can occur as a result of the feeding activities of insects or from disease. All of the foliage or the whole crown is affected.



>10%; <30% of Crown Affected—Cicada-caused flagging



>10%; <30% of Crown Affected—Gypsy moth feeding damage



>30% but not the Whole Crown—Oak wilt



Whole Crown Affected—Balsam woolly adelgid



Whole Crown Affected—Birch leafminer

# Signs and Symptoms of Branches and Bole

---

Insect and/or disease presence may be detected by careful inspection of the woody portions of the branches and bole. Boring insects produce frass and sawdust that can be seen in bark crevices or on the ground. Exit holes are a sign of insect activity. Signs of disease may include *fungus fruiting bodies* or *bleeding*.

## Insect Signs

Signs of insect-caused problems include frass or sawdust resulting from boring, or entrance or exit holes. These signs may be difficult to see without careful observation. Insects boring into pine trees often results in *pitch* or *sap* seen on the bole or branches.

### Frass Only

*Frass* is solid insect excrement usually consisting of a mixture of chewed plant fragments. Look for frass in bark crevices or at the base of the tree. Frass at the base of a tree may be a sign of insect feeding within the foliage.

### Sawdust

Wood-boring insects often produce what is essentially sawdust as they bore into the tree bole or branches. This sawdust can be found in bark crevices or at the base of the tree. For some *ambrosia beetles*, the sawdust appears almost like a toothpick sticking out from the tree bole.

### Pitch/Resin Exudation

Pitch or resin *exuding* from the bark is often seen on conifers as a result of insect activity, wounds, or disease. Don't confuse it with sapsucker damage. Sapsucker damage usually occurs in lines across or down the bole while pitch flowing from insect activity appears randomly across the bole.



Frass—Asian longhorned beetle



Frass—Pine webworm



Sawdust—Carpenter ants



Sawdust—Sawdust from boring of the striped Ambrosia beetle



Pitch/Resin Exudation—Red pine shoot moth



Pitch/Resin Exudation—Pitch tube from spruce engraver

# Signs and Symptoms of Branches and Bole

---

## **Insect Signs**

### **D-Shaped Exit Holes**

D-shaped exit holes are caused by a group of insects known as *buprestid* or *flat-headed beetles*. They produce oval to flattened tunnels within the bark and create a D-shaped exit hole as they exit the tree. The emerald ash borer and the oak splendour beetle both produce a D-shaped exit hole that is only about 6 mm across and can be difficult to see without careful observation of the tree bole. Trees with dieback should be examined for exit holes. Check ash and oak for D-shaped exit holes.

### **Pencil Round or Oval Exit Holes ( $\geq 2$ mm)**

Round exit holes in the branches or bole are created by insects as they leave the tree. Round-headed borers such as sawyers and two common ash borers—the banded ash borer and the red-headed ash borer—produce small, round exit holes. The Sirex woodwasp also produces a round exit hole (in pines) as does the Asian longhorned beetle (check maples carefully).

### **Shot Holes ( $< 2$ mm)**

Small beetles exiting a tree may leave tiny holes in the bark. A good example of this is small holes created by emerging ash bark beetles.

### **Other Holes**

There are many different types of holes made by insects in the branches and bole. If the holes found do not appear to fit into any of the above hole categories made by insects, please note here.



D-Shaped Exit Holes—Emerald ash borer



Pencil Round or Oval Exit Holes—Asian longhorned beetle



D-Shaped Holes—Bronze birch borer



Pencil Round or Oval Exit Holes—  
Sirex woodwasp exit holes



Shot Holes—Southern pine beetle  
exit holes

# Signs and Symptoms of Branches and Bole

---

## **Insect Presence**

Look for insects on the bark or the base of the tree. While caterpillars are most often found in the foliage, some may be crawling on the bark looking for a *pupation* site. Beetles may be searching for egg-laying sites. If you see ants, only the larger carpenter ants are of concern. Other signs of insects on the branches and bole include egg masses.

### **Caterpillars**

Caterpillars feed on foliage in the tree crown. Caterpillars found on tree branches and the bole are moving to a resting site or hiding.

### **Beetles**

Beetles can be found on the branches and bole. Females may be searching for egg-laying sites, or the adult beetle may be emerging from an exit hole.

### **Aphids**

Aphids are generally found on new growth and smaller twigs.

### **Scale**

Scales are generally found on the smaller branches and twigs, or on the bole of smooth-barked trees. A large scale infestation may exist before damage is obvious.



Caterpillars—Eastern tent caterpillar



Beetles—Asian longhorned beetle



Beetles—Adult emerald ash borer



Aphids—Hemlock woolly adelgid



Scale—Wax scale

# Signs and Symptoms of Branches and Bole

---

## **Insect Presence**

### **Carpenter Ants**

Carpenter ants make nests in wood by chewing sandpaper-smooth tunnels and chambers into the wood of trees. They feed on protein and sugar, mainly feeding on insects and sugars produced by aphids and scale insects. A common misconception is that carpenter ants eat wood. They do not eat wood but rather remove wood as they create galleries and tunnels, often impacting the structural integrity of trees and buildings. Wood is discarded as shredded fragments of coarse sawdust ejected from the nests. The sawdust may contain dead ants and bits of dead insects that the ants have eaten.

## **Disease Signs**

The structures produced by pathogens are referred to as disease signs. These can include *sporocarps* (mushrooms, conks, or other fruiting bodies), spores, and specialized vegetative and reproductive structures. Disease symptoms are the various responses by the host that are the result of the disease agent attacking the tree.

### **Decay**

Wood decay is a common occurrence among trees. Decay can affect the roots, sapwood, or heartwood of a tree. Some trees may appear to be healthy, yet have extensive decay. Trees with extensive decay are structurally weakened. Nearly all wood decay is caused by fungi. Mushrooms, sometimes called conks, are usually seen growing on the sides of trees. These are reproductive structures, commonly known as fungal fruiting bodies. Trees are usually infected for many years before mushrooms are visible.



Ants—Carpenter ants in cavity




Ants—Carpenter ant



Ants—Carpenter ant and associated feeding woodpecker damage



Decay—Decay cavity



Decay—Decay of branches and bole

# Signs and Symptoms of Branches and Bole

---

## Disease Signs

### Conks

**Conks** are spore-producing structures of some fungi that cause wood decay. Conks include structures that are hard and woody, or relatively soft and fleshy, as well as **bracket fungi** that are thinner and more flexible. Conks can be found individually, or in masses, but always grow on wood.

### Fleshy Mushrooms

Mushrooms are often prominent signs of plant disease and are some of the largest fungal fruiting bodies. Many mushrooms grow on wood and most that do cause wood decay. The extent of decay varies widely with each fungus.

### Cankers

A canker is a localized, often sunken, dead area on a twig, branch, or bole.

### Bleeding/Slime Flux

**Slime flux** is associated with several bacteria, and is an exudate that is composed of sap and bacteria. Slime flux often has a distinctive sharp odor of fermentation, and is the result of a bacterial infection of the **heartwood** area of the tree known as **bacterial wetwood**. Bleeding on branches and stems may be caused by insect attack as well as infection by fungi and bacteria. Bleeding on the stem is often the first sign of canker disease.



Conks—Artist's conk



Conks—Red-belted polypore (fungus)



Fleshy Mushrooms—Oyster mushroom



Cankers—Nectria canker



Cankers—Butternut canker



Bleeding/Slime Flux—Slime flux associated with bacterial wetwood

# Signs and Symptoms of Branches and Bole

---

## Disease Signs

### Resinosis/Gummosis

**Resinosis** is an accumulation of pitch in conifers that often arises as the result of insect or pathogen attack. Resinosis can occur anywhere on the woody portions of a tree. When the pitch from resinosis is present at the base of a tree, it often becomes mixed with soil, forming a hard, sticky substance that discolors the soil. **Gummosis** is the production of gummy substances on hardwoods, often in fruit trees, on the surface of the bark.

### Woody Galls or Burls

Woody galls or burls form on woody plants and are usually caused by bacteria invading the plant and affecting DNA. These galls can be found on tree trunks or on branches, and look like odd, corky balls. Pruning them away is the only cure.

Reference: Sustainable Urban Landscape Information Series (SULIS) [http://www.sustland.umn.edu/maint/tree\\_disease.html](http://www.sustland.umn.edu/maint/tree_disease.html)

## Problem Location

### Branches

Are the signs or symptoms of insects or disease found in the main branches?

### Bole and/or Root Collar

Are there symptoms or signs of insect and/or disease on the tree bole from the root collar up to the branches?



Resinosis/Gummosis—Gummosis caused by lesser peach tree borer



Resinosis—Sirex woodwasp boring symptom



Galls—Cynipid gall wasp



Gall—Crown gall on cherry bole



Burl—Burl and decay on pine bole



Disease Signs—Burl and decay on bole

# Signs and Symptoms of Branches and Bole

---

## Loose Bark

### Loose Bark

Loose or flaking bark may be a sign of serious problems within the tree. For example, a tree infested with bark beetles, wood-boring insects, or root and butt decay may have loose bark. Woodpecker feeding resulting in *bark sloughing* is a sign of an insect-infested tree. Sunscald and other physiological problems can also cause loose bark.

### Rhizomorphs Present

*Rhizomorphs* are shoestring-like structures produced by some root disease fungi. Rhizomorphs are usually dark-colored, tough structures that grow under the bark or in the soil. When they encounter a susceptible host, the rhizomorphs try to penetrate the bark and infect the host.

### Mycelial Fans or Pads Present

*Mycelial fans* or pads are wefts of tissue produced by fungi, usually found under the bark. Most are white or buff colored. Although mycelial fans are produced by several species of fungi, those produced by *Armillaria* spp. are probably most common.

### Insect Boring or Galleries causing Loose Bark

Boring insects attack both healthy and unhealthy trees. Larvae tunnel beneath the bark producing tunnels or galleries, which create entryways for pathogens that can cause structural weakness in the branches or bole. Woodborers can be detected in a number of ways: they can be heard feeding, frass and exit holes may be present, larvae may be exposed by peeling the bark, and feeding galleries and/or holes leading into the sapwood can be exposed by peeling back the bark. In severe infestations, the bark may be loose and peeling due to the galleries.



Loose Bark—*Armillaria* root rot



Rhizomorphs—Rhizomorphs of *Armillaria* root rot



Mycelial Fans—Mycelial fans of *Armillaria*



Insects Boring—Emerald ash borer



Insects Boring—Southern pine beetle galleries



# Glossary

---

**aphids**—small plant-eating/sucking insects that cause great damage to cultivated plants. Members of the order Homoptera that undergo simple metamorphosis and whose larvae somewhat resemble adults without wings

**abiotic stress**—impact on trees caused by non-living factors

**adventitious roots**—roots that arise at sites other than in the soil, such as roots originating on stems or leaves

**adventitious sprouts**—sprouts that develop on the trunk or branches that are often a result of poor pruning, injury, or boring insects

**ambrosia beetles**—beetles of the weevil subfamilies that live in symbiosis with ambrosia fungi and cultivate fungal communities that provide their sole source of nutrition

**bacteria**—a large group of unicellular microorganisms that have a wide range of shapes and are capable of living in all organic lifeforms

**bacterial wetwood**—a common disease that affects the wood and bark of trees, characterized by water-soaked wood and a liquid that bleeds from wounds and becomes slimy, resulting in a condition known as slime flux

**bark sloughing**—the casting off of bark related to tree disease or stress

**biotic**—relating to, produced by, or caused by living organisms

**bleeding**—the production of sap/fluid in response to disease, insects, or wounding

**bole**—the trunk of a tree

**bracket fungi**—shelf fungi that produce shelf or bracket-shaped fruiting bodies (conks) and range from a single row of a few to dozens of caps

**buprestid**—a family of beetles with a long, flat, metallic-colored body and larvae that are harmful to woody plants

**cambium**—the cambium layer is found directly beneath the bark on trees and is surrounded on the outside by phloem and inside by xylem

**canker**—a localized, often sunken, dead area on a twig, branch, or bole

**causal factor**—anything that causes a disruption of a plant's normal growth

**chlorosis**—complete yellowing of leaves

**chlorotic**—plant tissue that appears pale green to yellow

**conks**—spore-producing structures of some fungi that cause wood decay

**defoliation**—foliage eaten wholly or partially

**deicing salts**—salts used for ice and snow removal that are harmful to trees and result in abnormal foliage color, needle tip burn, and marginal leaf burn

**drifts**—when herbicides or pesticides are moved by wind or other factors from an intended target to a nearby nonintended target

**epicormic sprouts**—shoots that arise from latent or adventitious buds that occur on stems and branches or from suckers at the base of the tree

**exit holes**—the holes left by young adults as they leave the tree after the larvae of insects pupate and adults hatch

**exude**—to discharge slowly

**flagging**—the yellowing or wilting of foliage on a branch that commonly occurs with a number of tree diseases

**flat-headed beetles**—beetles in the family Buprestidae that are beautifully marked, metallic colored, vary in size, and are somewhat flattened and boat shaped. Although they are beetles, they are referred to as borers because of the damage they cause in trees

**fleshy mushroom**—the spore-bearing fruiting body of a fungus typically produced above ground on soil or on its food source

**flush cut**—pruning cuts that originate inside the branch bark ridge or the branch collar, causing unnecessary injury to stem tissue

**frass**—solid insect excrement usually consisting of a mixture of chewed plant fragments

**frost cracks**—cracks that are caused by differential expansion and contraction of woody stem tissues during extremely cold periods

**fungus fruiting body**—a fungal structure that produces spores (see “sporocarp”)

**galls**—plant tissue that has developed as the result of feeding or other activity of insects or mites

**galleries**—series of tunnels created beneath the bark in the cambium by wood boring insects

**gummosis**—an accumulation of pitch in conifers that is often the result of insect or pathogen activity

**heartwood**—nonliving xylem cells that form a core of wood in the center of the stem that provides the structural strength in trees

**herbicide**—a chemical substance or cultured biological organism used to kill or suppress the growth of plants

**honeydew**—a sticky, sweet substance that is a good substrate for the growth of several different fungi that have very dark mycelium

**interveinal chlorosis**—yellowish discoloration of green tissues in areas between the leaf veins that is usually related to lack of chlorophyll production due to disease or stress

**interveinal scorching**—the discoloration in the tissues between the leaf veins

**leaf mining**—the serpentine pattern of feeding beneath the leaf surface as insects live or feed within a leaf

**leaf rollers**—insects that construct shelters by rolling leaves with silk webbing

**marginal scorching**—dead tissues on the margins of leaves that result in browning and shriveling of foliage

**midrib**—the midvein of a leaf or leaflet

**mottling**—spotting of leaves or needles in an uneven and undefined pattern, usually involving several different shades of color

**mycelial fans**—wefts of tissue produced by fungi, usually occurring just under the bark

**mycelium**—mycelium (plural mycelia) is the vegetative part of a fungus, consisting of a mass of branching, thread-like structures

**necrotic leaf blotches**—varying degrees of dead areas or spots on leaves, often used to describe brown spots left by insects or disease

**nontarget**—not the intended target; term related to chemical applications when drift from herbicides/insecticides/fungicides causes damage to adjacent nontarget vegetation

**parasitic plant**—derives some or all of its sustenance from another plant. Parasitic plants have a modified root that penetrates the host plant and connects to the xylem, phloem, or both

**pathogens**—biological agents that cause disease or illness to their host. Soil contamination has the longest or most persistent potential for harboring a pathogen.

**pervasive twig dieback**—twig dieback that spreads throughout every part of the tree

**pitch**—a highly viscous liquid that appears solid; also known as resin

**phloem**—in vascular plants, phloem is the living tissue that carries organic nutrients, particularly sucrose, a sugar, to all parts of the plant

**pupation**—period in the life cycle of an insect when the adult structures are formed and the larval structures are broken down

**pustule**—an abscess or eruption

**resinosis**—exudations of pitch on the tree, often associated with fungal infection of the roots, bole, or branches, or with root collar weevils

**rhizomorphs**—shoestring-like structures produced by some root disease fungi

**sap**—phloem and xylem fluid released from a tree due to the buildup of pressure in sapwood that is released through wounds or openings

**sapsuckers**—birds within the woodpecker family that drill holes in the bark of trees while looking for insects to feed upon

**sawflies**—sawflies belong to the same group of insects as wasps, ants, and bees

**scorching**—dead tissues on the margins of leaves or between veins that result in browning and shriveling of foliage

**shot holes**—tiny holes found in the bark caused by beetles exiting a tree

**sign**—the physical evidence of a causal agent

**skeletonizing**—the lacy effect on leaves created by insects feeding on only the softer parts of leaves

**slime flux**—slime flux is associated with several bacteria and is an exudate that is composed of sap and bacteria

**spindle-shaped bags**—small, spindle-shaped bags hanging from the foliage or twigs that are the protective feeding or reproductive enclosures for some caterpillars

**sporadic feeding**—feeding by insects that occurs irregularly or at intervals that have no apparent pattern but can cause significant injury to plants

**sporocarp**—a specialized structure with the primary function of producing and releasing spores.

**stippling**—small, white-spotted, speckled injuries on foliage caused by certain insects, mites, and viruses. In some cases it may be yellow or black

**symptom**—a plant's reaction to a causal agent

**webworm**—a caterpillar that spins a web in which it feeds or rests

**wilt**—the loss of water turgor pressure in a leaf, causing a leaf to droop, curl, or lose a degree of its normal color

**witches' brooms**—deformities in a tree that change the natural structure of branches or twigs. They often arise as the result of infection by mistletoes, fungi, insects, or disease

**wefts**—threadlike or weaved as often seen with mycelial fans, usually occurring under the bark

# Other Resources

---

## Project Links:

IPED Protocol Home Page <http://pest.itreetools.org/>

IPED Online Resources <http://wiki.bugwood.org/IPED>

## Resources:

U.S. Forest Service, Forest Health Protection, Northeastern Area State and Private Forestry,  
<http://www.na.fs.fed.us/fhp/index.shtm>

U.S. Forest Service, Forest Pest Alerts [www.na.fs.fed.us/pubs/palerts.shtm](http://www.na.fs.fed.us/pubs/palerts.shtm)

National Plant Diagnostic Network (Diagnostic Laboratory, First Detector Training)  
[www.npdn.org/DesktopDefault.aspx](http://www.npdn.org/DesktopDefault.aspx)

APHIS – Animal and Plant Health Inspection Service <http://www.aphis.usda.gov/>

i-Tree Tools for Assessing and Managing Community Forests [www.itreetools.org/](http://www.itreetools.org/)

Forestry Images: Forest Health, Natural Resources, and Silviculture Images  
<http://www.forestryimages.org/>

## References:

Lloyd, John [coordinating author]; Kruidenier, Bill; Moell, Todd; Skiera, James. 2007. Plant health care for woody ornamentals: a professional's guide to preventing and managing environmental stresses and pests. University of Illinois, Office of Agricultura. 256 p.

Sinclair, Wayne A.; Lyon, Howard H. 2005. Diseases of trees and shrubs. 2d ed. Cornell University Press. 660 p.

# Photo Credits

---

- 0014295 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 0355012 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 0364092 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 0364100 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 0907049 Photo by E. Bradford Walker, Vermont Department of Forests, Parks and Recreation, Bugwood.org
- 1141006 Photo by Gyorgy Csoka, Hungary Forest Research Institute, Bugwood.org
- 1193010 Photo by Dean Morewood, Health Canada, Bugwood.org
- 1210083 Photo by Dave Powell, USDA Forest Service, Bugwood.org
- 1234182 Photo by Clemson University - USDA Cooperative Extension Slide Series, Bugwood.org
- 1238068 Photo by Randy Cyr, Greentree, Bugwood.org
- 1238102 Photo by Randy Cyr, Greentree, Bugwood.org
- 1238111 Photo by Randy Cyr, Greentree, Bugwood.org
- 1241596 Photo by Scott Tunnock, USDA Forest Service, Bugwood.org
- 1241745 Photo by David Beckman, Idaho Department of Lands, Bugwood.org
- 1241757 Photo by Susan K. Hagle, USDA Forest Service, Bugwood.org
- 1254083 Photo by William M. Ciesla, Forest Health Management International, Bugwood.org
- 1263084 Photo by E.G. Kuhlman, USDA Forest Service, Bugwood.org
- 1296183 Photo by Gyorgy Csoka, Hungary Forest Research Institute, Bugwood.org
- 1301043 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org

1304003 Photo by Carroll E. Younce, USDA Agricultural Research Service, Bugwood.org

1371075 Photo by Andrej Kunca, National Forest Centre - Slovakia, Bugwood.org

1393012 Photo by Dennis Haugen, USDA Forest Service, Bugwood.org

1393019 Photo by Dennis Haugen, USDA Forest Service, Bugwood.org

1393029 Photo by Dennis Haugen, USDA Forest Service, Bugwood.org

1397126 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org

1398035 Photo by Steven Katovich, USDA Forest Service, Bugwood.org

1398266 Photo by Steven Katovich, USDA Forest Service, Bugwood.org

1406224 Photo by USDA Forest Service - North Central Research Station Archive, USDA Forest Service, Bugwood.org

1406274 USDA Forest Service - North Central Research Station Archive, USDA Forest Service, Bugwood.org

1427015 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org

1439005 Photo by Edward Czerwinski, Ontario Ministry of Natural Resources, Bugwood.org

1457027 Photo by Steven Katovich, USDA Forest Service, Bugwood.org

1458059 Photo by North Carolina Forest Service Archive, Bugwood.org

1460031 Photo by James W. Smith, USDA APHIS PPQ, Bugwood.org

1467013 Photo by USDA Forest Service - Ogden Archive, USDA Forest Service, Bugwood.org

1473007 Photo by Rocky Mountain Research Station/Forest Pathology Archive, USDA Forest Service, Bugwood.org

1473055 Photo by Rocky Mountain Research Station/Forest Pathology Archive, USDA Forest Service, Bugwood.org

1492062 Photo by University of Georgia Plant Pathology Archive, University of Georgia, Bugwood.org

- 1505056 Photo by Paul A. Mistretta, USDA Forest Service, Bugwood.org
- 1626029 Photo by John A. Weidhass, Virginia Polytechnic Institute and State University, Bugwood.org
- 1748005 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 1949012 Photo by C.E. Seliskar, Bugwood.org
- 1949013 Photo by C.E. Seliskar, Bugwood.org
- 2101035 Photo by Daniel Adam, Office National des Forêts, Bugwood.org
- 2108027 Photo by Ronald F. Billings, Texas Forest Service, Bugwood.org
- 2108041 Photo by Ronald F. Billings, Texas Forest Service, Bugwood.org
- 2121089 Photo by Steven Katovich, USDA Forest Service, Bugwood.org
- 2200041 Photo by Whitney Cranshaw, Colorado State University, Bugwood.org
- 2251099 Photo by USDA Forest Service Archive, USDA Forest Service, Bugwood.org
- 3036022 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 3036037 Photo by Robert L. Anderson, USDA Forest Service, Bugwood.org
- 3046072 Photo by Theodor D. Leininger, USDA Forest Service, Bugwood.org
- 3067038 Photo by James Solomon, USDA Forest Service, Bugwood.org
- 3225056 Photo by Ronald F. Billings, Texas Forest Service, Bugwood.org
- 3226030 Photo by Ronald F. Billings, Texas Forest Service, Bugwood.org
- 3570012 Photo by Petr Kapitola, State Phytosanitary Administration, Bugwood.org
- 4212011 Photo by Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org
- 4212014 Photo by Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org

- 4212031 Photo by Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org
- 4212056 Photo by Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org
- 4213094 Photo by Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org
- 4215048 Photo by Fred Baker, Utah State University, Bugwood.org
- 4798040 Photo by Kenneth R. Law, USDA APHIS PPQ, Bugwood.org
- 4836011 Photo by E. Bradford Walker, Vermont Department of Forests, Parks and Recreation, Bugwood.org
- 5016094 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5016097 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5018075 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5019024 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5020025 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5020027 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5020085 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5021069 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org
- 5032082 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5032085 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org

- 5033003 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5034082 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5036036 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5038017 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5038047 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5038050 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5039050 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5047087 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5049050 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5051049 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5052017 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5052043 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5052096 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5054052 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5055016 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5056022 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5059018 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5083042 Photo by Whitney Cranshaw, Colorado State University, Bugwood.org
- 5123062 United States National Collection of Scale Insects Photographs Archive, USDA Agricultural Research Service, Bugwood.org
- 5131099 Photo by Pennsylvania Department of Conservation and Natural Resources - Forestry Archive, Bugwood.org

- 5137091 Photo by Susan Ellis, Bugs in the System, Bugwood.org
- 5179072 Photo by Jeffrey Eickwort, Florida Department of Agriculture and Consumer Services, Bugwood.org
- 5202059 Photo by Steven Katovich, USDA Forest Service, Bugwood.org
- 5291028 Photo by Joseph LaForest, University of Georgia, Bugwood.org
- 5291065 Photo by Joseph LaForest, University of Georgia, Bugwood.org
- 5298042 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5298046 Photo by Joseph O'Brien, USDA Forest Service, Bugwood.org
- 5360705 Photo by Eugene E. Nelson, Bugwood.org
- 5365410 Photo by H.J. Larsen, Bugwood.org
- 5367164 Photo by William Jacobi, Colorado State University, Bugwood.org
- 5368719 Photo by Paul Bachi, University of Kentucky Research & Education Center, Bugwood.org
- 5369161 Photo by Marianne Prue, Ohio Department of Natural Resources - Division of Forestry, Bugwood.org
- 9009023 Photo by Ronald F. Billings, Texas Forest Service, Bugwood.org

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

